

## NEWS

# ENHANCED RECOVERY AFTER ARTHROPLASTY: A TEAM EFFORT

This article will outline the collaborative approach undertaken between anaesthetic and orthopaedic specialists at Vermont Private Hospital in Victoria to facilitate the successful undertaking of a trial program of enhanced recovery after total joint arthroplasty. The rationale of this approach is that short-stay or outpatient arthroplasty is an emerging model of care that should not be a goal in itself, but rather be the result of specialised multidisciplinary coordination implementing a holistic program that can optimise patient recovery. For enhanced recovery to be achieved in this model, an investment in time and resources is required by healthcare providers to deliver and support a unique skillset and willingness to collaborate closely with relevant team members involved.

Total hip and knee arthroplasty are common orthopaedic procedures, which are continuing to increase in frequency in the setting of rising rates of obesity and osteoarthritis, and an aging Australian population which desires to maintain their functionality.<sup>1,2</sup> Arthroplasty procedures have typically been associated with an average hospital length of stay of 5.4 +/- 1 day duration after which 43% of patients proceed to ongoing inpatient rehabilitation.<sup>3</sup>

Several overseas countries have well-established fast track arthroplasty programs in which patients undergoing elective hip and knee replacements remain in hospital for 1-2 days, and in some instances less than 24 hours, with

the bulk of the recovery process occurring in the outpatient setting.<sup>4,5,6</sup> Until recently this approach has not been commonly practiced in Australia due to factors that include lack of familiarity with the process by healthcare providers, particularly in regards to patient safety and outcomes, private health insurance reimbursement criteria, competing interests of specialists and a lack of coordinated processes between the hospital and specialists. To challenge these problems and overcome the barriers to implementation there needs to be a supportive healthcare provider environment, a well-coordinated and specialised multidisciplinary team in place and private health insurance involvement.

When patient care interventions operate in isolation it is unlikely to lead to beneficial healthcare value or improved outcomes for patients. Enhanced recovery protocols are bundled interventions of integrated multidisciplinary care that allows patients to benefit from a variety of evidence-based care targeted at a shared objective – global modulation of the surgical stress response and early mobilisation of the joint to facilitate improved range of movement and patient outcomes.<sup>7</sup> Meta-analyses indicate that patients have the potential to benefit significantly from enhanced recovery after arthroplasty programs.<sup>8</sup> In addition to the psychological benefit and satisfaction of recovering in the home rather than the hospital setting, patients have been shown to experience reduced mortality, reduced morbidity in the form of thromboembolic events, cardiopulmonary complications,

delirium and sleep disturbance, and better analgesia with less pain and joint stiffness, all without increasing the risk of readmission or adverse events.<sup>9-13</sup>

The ERAS® Society suggests that there are many components of care that go into the implementation of 'enhanced recovery programs' and as a result the development of our rapid recovery after elective hip and knee arthroplasty trial pathway at Vermont Private Hospital is a multidisciplinary endeavour involving the orthopaedic surgeon, anaesthetist, orthopaedic nurse practitioner, physician, physiotherapists and nursing staff. The anaesthetist has a vital role to play in every stage, from pre-admission to well after discharge, to ensure that patients are able to progress smoothly through this 'novel' model of care and thus facilitate a significant opportunity for patient-centric gains. Whilst an in-depth discussion of every component is beyond the scope of this article, the generic framework is divided into the key three phases of the patient journey: pre-operatively, intra-operatively and post-operatively. The highlights of the trial pathway are as follows:

Pre-operatively:

- Patient selection: Patients are assessed for their appropriateness for this model of care based on the combination of results from an anaesthetic pre-assessment including a risk calculation, and an orthopaedic pre-operative survey constructed to predict discharge disposition after total joint arthroplasty known as RAPT.

- **Medical optimisation:** This comprehensive screening allows for early physician intervention to address modifiable risk factors and also allows post-operative planning for management of medical co-morbidities in an outpatient setting.
- **Patient education:** All patients are encouraged to attend a 'joint school' run by an orthopaedic nurse practitioner, where they are taught what to expect at each stage, given advice about the surgical and anaesthetic plan (type of anaesthesia, fasting/diet/aperients and the analgesia plan from PACU to post-discharge), and are provided with hands-on instruction about physiotherapy and post-operative mobilisation. The key aim is active participation of the patient in their own recovery and setting of appropriate expectations.
- **Pre-operative fasting/carbohydrate loading:** There is limited data in regards to metabolic state and recovery after joint arthroplasty, but given the evidence in other surgery our aim is for well-hydrated patients in a 'fed' state.<sup>14</sup>

#### Intra-operatively:

The orthopaedic surgeon needs to collaborate closely with the anaesthetist to facilitate a plan that will modulate the surgical stress response and allow early mobilisation. This cooperation is a critical component of enhanced recovery after surgery pathways.

- **Anaesthetic technique:** In our protocol we prefer a 'low dose' (5-7.5mg bupivacaine) neuraxial technique. Evidence indicates that this provides the optimal attenuation of the surgical stress response and acts as a bridge to early post-operative analgesia, significantly reducing the risk of moderate-to-severe postoperative pain, 15 whilst still allowing resolution of the motor block and ambulation within  $\leq 300$  minutes in 95% of patients. This rapid improvement facilitates physiotherapy intervention from PACU onwards, with improved mobility and rehabilitation.<sup>16,17</sup> There is also evidence

that it may even reduce in-hospital complications and 30-day mortality.<sup>18,19</sup>

- **Anaesthetic adjuncts:** There are a number of controversial areas that require further research to clearly elucidate the best outcome for arthroplasty patients, and our pathway will evolve with the results from continued research. Based on currently available evidence we use the following adjuncts: maintaining normothermia,<sup>20,21</sup> prevention of hypertensive anaesthesia with the use of neuraxial anaesthesia, blood conservation strategies with the use of tranexamic acid and cell salvage,<sup>22,23</sup> TIVA and opioid minimisation for sedation/GA,<sup>24</sup> goal-directed fluid therapy with early return of oral intake,<sup>25</sup> weight adjusted cephazolin as anti-microbial prophylaxis to target adequate joint concentration,<sup>26,27</sup> pre-operative pregabalin to reduce opioid consumption and neuropathic pain,<sup>28,29</sup> and peri-operative systemic steroid.<sup>30</sup>
- **Surgical interventions:** Minimally invasive surgical techniques such as anterior approach hip replacement are usually

required to facilitate early discharge. In knee replacement surgery, minimising tourniquet time and using a kinematic alignment protocol to reduce soft tissue trauma may assist in rapid recovery.

#### Post-operatively:

- **Opioid-sparing multimodal analgesia:** Paracetamol and NSAID's are the backbone of the analgesia regimen. With respect to advanced analgesic modalities, intrathecal opiates +/- PCEA, peripheral nerve block and peri-articular injection all appear reasonable options. The challenge is that there are few studies which directly compare these modalities, although a recent meta-analysis by Xing et al has demonstrated that there is a synergistic effect between LIA and ACB which resulted in a statistically significant improvement in analgesia and lowered opioid consumption when these two modes of anaesthesia were used in combination rather than individually.<sup>31,32</sup> As such, this is the approach we have implemented within our protocol. Continuous peripheral nerve block



# NEWS

provides the same efficacy of analgesia as PCEA,<sup>33</sup> allowing reduced subsequent opioid consumption, but with decrease in nausea and vomiting, improvement in mobilisation, reduction in the ongoing surgical stress response and potentially a reduction in morbidity and mortality. With appropriate pre-operative education and follow-up patients can be discharged home with their CPNB in-situ, with no evidence of increased risk of falls,<sup>34,35</sup> thus allowing the same analgesia they would receive within the hospital setting.

- Team-effort and follow-up: From admission to discharge everyone involved in the patient's care needs to be educated on the components of the pathway. Once patients are settled into their own environment at home it is imperative that they have clear guidelines on how to access extra assistance if required, such that there recovery is enhanced rather than hindered.

In order to change the current approach to enhanced recovery after joint arthroplasty, development of process, practice and communication is required. This includes a multidisciplinary team, of which the anaesthetic practices outlined above are integral component. There must be a holistic and targeted approach that works in a coordinated fashion to support improved outcomes for the patient, and as a result facilitate an early discharge from hospital. Overall the implementation of such an approach aims to challenge the status quo model for elective hip and knee procedures with Australasia, but the early results from the implementation of this collaborative approach between orthopaedic surgeon, anaesthetist, nursing staff, allied health practitioners, hospital and health insurer highlights the potential to facilitate early discharge, with improved patient experience and outcomes, without compromising safety.

Mr Daevyd Rodda  
Dr Dwane Jackson

## References

1. AIHW analysis of National Hospital Morbidity Database 2014-15 and ABS Estimated Resident Population, 30 June 2014
2. Australian Orthopaedic Association. National Joint Replacement Registry annual report 2015. Adelaide: AOA; 2015.
3. Schilling C, Keating C, Barker A, Wilson S, Petrie D. 'Predictors of inpatient rehabilitation after total knee replacement: an analysis of private hospital claims data', *Med J Aust* 2018; 209(5):222-227.
4. Xu J, Cao, J, Chaggar G, Negus J. 'Comparison of outpatient versus inpatient total hip and knee arthroplasty: A systematic review and meta-analysis of complications', *J Orthop* 2019; 13(17):38-43.
5. Vehmeijer S, Husted H, Kehlet, H. 'Outpatient total hip and knee arthroplasty: Facts and challenges', *Acta Orthopaedica* 2017; 89(2):1-4.
6. Krause A, Sayeed Z, El-Othmani M, Palakkonda V, Mihalko W, Saleh K. 'Outpatient total knee arthroplasty: Are we there yet? (Part 1&2)', *Orthop Clin North Am* 2018; 49(1):1-6, 7-16.
7. Soffin E, YaDeau J. 'Enhanced recovery after surgery for primary hip and knee arthroplasty: A review of the evidence', *Br J Anaesth* 2016; 117(3):iii62-iii72.
8. Deng Q, Gu H, Peng W, Zhang Q, Huang Z, Zhang C, Yu Y. 'Impact of enhanced recovery after surgery on postoperative recovery after joint arthroplasty: results from a systematic review and meta-analysis', *Postgrad Med J* 2018; 94(1118): 678-693.
9. Vendittoli P, Pellei K, Desmeules F, Masse V, Loubert C, Lavigne M, Fafard J, Fortier L. 'Enhanced recovery short-stay hip and knee joint replacement program improves patients outcomes while reducing hospital costs', *Orthop Traumatol Surg Res* 2019; 105(7):1237-1243.
10. Barbieri AVanhaecht KVan Herck P et al. 'Effects of clinical pathways in the joint replacement: a meta-analysis', *BMC Med* 2009; 7:32.
11. Malviya A, Martin K, Harper I et al. 'Enhanced recovery program for hip and knee replacement reduces death rate. A study of 4500 consecutive primary hip and knee replacement', *Acta Orthop* 2011; 82:577-581.
12. Lovald S, Ong K, Malkani A, Lau E, Jordana K, Schmier M, Kurtz S, Manley M. 'Complications, mortality, and costs for outpatient and short-stay total knee arthroplasty patient in comparison to standard-stay patients', *J Arthro* 2014; 29(3): 510-515.
13. Berg U, BuLow E, Sundberg M, Rolfson O. 'No increase in readmission or adverse events after implementation of fast-track program in total hip and knee replacement at 8 Swedish hospitals: An observational before-and-after study of 14,148 total joint replacements 2011-2015', *Acta Orthopaedica* 2018; 89(5):522-527.
14. Alito M, Ade Aguilar-Nascimento J. 'Multimodal perioperative care plus immunonutrition versus traditional care in total hip arthroplasty: a randomized pilot study', *Nutr J* 2016; 15:34.
15. Liu S, Buvanendran A, Rathmell JP et al. 'Predictors for moderate to severe acute postoperative pain after total hip and knee replacement', *Int Orthop* 2012; 36:2261-2267.
16. Awad I, Cheung J, Al-Allaq Y et al. 'Low-dose spinal bupivacaine for total knee arthroplasty facilitates recovery room discharge: a randomized controlled trial', *Can J Anesth* 2013; 60:259-265.
17. Lemoine A et al. 'Modelling of the optimal bupivacaine dose for spinal anesthesia in ambulatory surgery based on data from systematic review', *Eur J Anaesthesiol* 2016.
18. Memtsoudis S, Sun X, Chiu Y, Studner O, Liu S, Banerjee S, Mazumdar M, Sharrock N. 'Perioperative comparative effectiveness of anaesthetic technique in orthopaedic patients', *Anaesthesiology* 2013; 118(5):1046-1058.
19. Perlas A, Chan V, Beattie S. 'Anesthesia technique and mortality after total hip and knee arthroplasty: A retrospective, propensity score-matched cohort study', *Anesthesiology* 2016; 125(4):724-731.
20. Winkler M, Akça O, Birkenberg B et al. 'Aggressive warming reduces blood loss during hip arthroplasty', *Anesth Analg* 2000; 91:978-984.
21. Benson E, McMillan D, Ong B. 'The effects of active warming on patient temperature and pain after total knee arthroplasty', *Am J Nurs* 2012; 112: 26-33.
22. Oremus K, Sostaric S, Trkulja V, Haspl M. 'Influence of tranexamic acid on postoperative autologous blood retransfusion in primary total hip and knee arthroplasty: a randomized controlled trial', *Transfusion* 2014; 54:31-41.
23. Wei Z, Liu M. 'The effectiveness and safety of tranexamic acid in total hip or knee arthroplasty: a meta-analysis of 2720 cases', *Transfus Med* 2015; 25:151-162.
24. Harsten A, Kehlet H, Ljung P, Toksvig-Larsen S. 'Total intravenous general anaesthesia vs spinal anaesthesia for total hip arthroplasty: a randomised controlled trial', *Acta Anaesthesiol Scand* 2015; 59:298-309.
25. Benes JHaidingerova LPouska J et al. 'Fluid management guided by a continuous non-invasive arterial pressure device is associated with decreased postoperative morbidity after total knee and hip replacement', *BMC Anesthesiol* 2015; 15: 148.
26. Voigt J, Mosier M, Darouiche R. 'Systematic review and meta-analysis of randomized controlled trials of antibiotics and antiseptics for preventing infection in people receiving primary total hip and knee prostheses', *Antimicrob Agents Chemother* 2015; 59:6696-6707.
27. Sanders F, Gosling J, Mathot R, Scheoers T. 'Target site antibiotic concentrations in orthopedic/trauma extremity surgery: is prophylactic cefazolin

- adequately dosed? A systematic review and meta-analysis', *Acta Orthop* 2019; 90(2):97-104.
28. Mao Y, Wu L, Ding W. 'The efficacy of preoperative administration of gabapentin/pregabalin in improving pain after total hip arthroplasty: a meta-analysis', *BMC Musculosk Disord* 2016; 17:373.
  29. Chao H, Ming-Jie K, Jian-Xiong M, Xin-Long M. 'Is pregabalin effective and safe in total knee arthroplasty? A PRISMA-compliant meta-analysis of randomized controlled trials', *Medicine* (Baltimore). 2017 96(26):e6947.
  30. Yue C, Wei R, Liu Y. 'Perioperative systemic steroid for rapid recovery in total knee and hip arthroplasty: a systematic review and meta-analysis of randomized trials', *J Orthop Surg Res* 2017; 12(1):100.
  31. Xing Q, Dai W, Zhao D, Wu J, Huang C, Zhao Y. 'Adductor canal block with local infiltrative analgesia compared with local infiltrate analgesia for pain control after total knee arthroplasty: A meta-analysis of randomized controlled trials', *Medicine* (Baltimore). 2017; 96(38):e8103.
  32. Sawhney M, Mehdian H, Kashin B, Ip G, Bent M, Choy J, McPherson M, Bowry R. 'Pain after unilateral total knee arthroplasty: A prospective randomized controlled trial examining the analgesic effectiveness of a combined adductor canal peripheral nerve block with periarticular infiltration versus adductor canal nerve block alone versus periarticular infiltration alone', *Anesth Analg* 2016;122(6):2040-2046.
  33. Paul J, Arya A, Hurlburt L, Cheng J, Thabane L, Tidy A, Murthy Y. 'Femoral nerve block improves analgesia outcomes after total knee arthroplasty: a meta-analysis of randomized controlled trials', *Anesthesiology* 2010; 113(5):1144-1162.
  34. Seung S, Gul K, Jin H, Do H, Youn G, Beyoung Y. 'Comparison of the effect of continuous femoral nerve block and adductor canal block after primary total knee arthroplasty', *Clin Orthop Surg* 2017; 9(3):303-309.
  35. Yugal K, Ramneek M, Abhimanyu K, Amol P, Mukul C. 'A comparative analysis of femoral nerve block with adductor canal block following total knee arthroplasty: A systematic literature review', *J Anaesthesiol Clin Pharmacol* 2018; 34(4):433-438.

## ABOUT THE AUTHORS

**Mr Daevyd Rodda** is an orthopaedic surgeon with extensive experience in hip and knee arthroplasty. He has a special interest in rapid recovery surgical techniques, and teaches these techniques to surgeons around Australia and internationally. Mr Rodda established the rapid recovery joint replacement program at Vermont Private Hospital and is a strong advocate of value-based health care in orthopaedic surgery.

**Dr Dwane Jackson** is a consultant anaesthetist with significant experience in arthroplasty surgery. He is actively involved in research, having published articles in peer-reviewed journals and has several research projects underway. With a special area of interest in holistic perioperative medicine utilising enhanced recovery after surgery techniques he has been a key driver in implementing the latest evidence-based techniques into various specialities at both tertiary public institutions and privately such as this current orthopaedic project.